Product Manual



Protease Inhibitor Cocktail (Multipurpose), EDTA (Combined)

Catalog# BWR1094

Size: 1 set

Lot # Check on the product label

Introduction

This product provides complete protection of proteins from degradation by endogenous proteases that may be released during extraction and purification of proteins from animal and plant tissues, cultured cells, yeast or bacteria. While extraction of proteins, tissue or cell lysis buffer is filled with enzyme which produced by lysosome, and the unactive proteases are fully activated, the latter begin to digest their surrounding proteins. To avoid this, the protease inhibitor must be added into the lysis buffer to prevent the digestion of proteases. The compositions of proteases that come from animal (mammal), plant, mould or bacterial are quite different, and will be changed based on the development of the organism state. So, it's important to choose the right protease inhibitor while extraction and electrophoresis of various kinds of proteins. This product is a mixture of 6 different protease inhibitors, this ensures inhibition of a broad range of proteases and esterases, including aspartase, cysteine, serine and aminopeptidase.

Application: Mammalian cell and tissue extracts.

Specificity: Serine protease, Cysteine protease, Insulin-like protease, Aspartase, and Aminopeptidase.

Appearance: Liquid in 1ml of DMSO.

Kit Components

Components	Size	Storage Instruction
Protease Inhibitor Cocktails (Multipurpose)	1ml (100×)	Store at -20℃ for one year.
EDTA	1ml (100×)	Store at -20℃ for one year.

Instruction

Dilute the two components at 1:100 in an appropriate lysis buffer and mix thoroughly, then extract the sample protein, or according to the purpose of experiment, and the cultured cells and tissues, grope and optimize the experimental conditions through experiment.